



Key references

- The Action to Control Cardiovascular Risk in Diabetes Study Group. Effects of Intensive glucose lowering in type 2 diabetes. *N Engl J Med* 2008;358:2545-59
- The ADVANCE Collaborative Group. Intensive blood glucose control and vascular outcomes in patients with type 2 diabetes. *N Engl J Med* 2008;358:2560-72
- Duckworth W, et al. Glucose control and vascular complications in veterans with type 2 diabetes. *N Engl J Med* 2009;360.
- Herman GA, et al. Pharmacokinetics and pharmacodynamics of sitagliptin, an inhibitor of dipeptidyl peptidase IV, in healthy subjects: results from two randomized, double-blind, placebo-controlled studies with single oral doses. *Clinical pharmacology and therapeutics*. 2005. 78(6):675-88.
- Kendall DM et al. Effects of exenatide on glycaemic control over 30 weeks in patients with type 2 diabetes treated with metformin and a sulphonylurea. *Diabetes care*. 2005. 28:1083-1091

- Nauck et al. LEAD-2: Liraglutide added on to metformin versus glimepiride. *Diabetes Care*. 2009. 32:84-90
- Garber et al. LEAD-3: Liraglutide as monotherapy versus glimepiride. *Lancet*. 2009. 373:473-481